



1<sup>st</sup> International Symposium  
on  
***Nanomedicine: Past, Present & Future Prospects***  
December 20-21, 2010  
&  
Workshop  
on  
***Techniques in Nanomedicine Research***  
December 22-23, 2010

**Anti-Emetic Activity of Six Aromatic Medicinal Plants by Using Chick Emetic Model**

**Salman Ahmed**, Iqbal Azhar, M.Mohtasheemul Hasan and Syed Waseemuddin Ahmed

Department of Pharmacognosy, Faculty of Pharmacy, University of Karachi, Karachi – 75270, Pakistan. mmhr260@hotmail.com

Current study was conducted to explore the anti-emetic activity of six aromatic medicinal plants viz *Carissa carandus* L., *Cichorium intybus* L., *Cinnamomum tamala* L., *Lallemantia royleana* Benth., *Matricaria chamomilla* L., and *Piper metysticum* G. Forst. Anti-emetic activity was assessed using chick as anti-emetic model. The ethanol extracts of these plants were administered at 150 mg/kg BW orally. *Lallemantia royleana* and *Piper metysticum* showed significant ( $P < 0.001$ ) decrease in retches induced by copper sulphate pentahydrate given orally at 50 mg/kg BW. Domperidone was given at 100 mg/kg as a reference drug.

**References.**

1. Yang, Y., Kinoshita, K., Koyania, K., Takahashi, K., Tai, T., Nuroura Y. and Watanabe, K. *Phytomed.* **6**(2): 89-93 (1999).
2. Parvez, H. S., Minami, M., Caudy P., Endo, T., Parvez, S., Hirafuji, M., Hamaue, N., Akita, K., Saito, H., Yoshioka, M., and Qureshi, G. *Neuroendocrinol. Lett.*, **18**(2/3): 85-102 (1998).
3. Nimri, L.F., Meqdam, M.M. and Alkofahi, A. *Pharm. Bio.*, **37**(3):196-201, (1999).