

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/289857944>

Total, insoluble and soluble dietary fiber contents of three Pakistani legume flours

POSTER · JANUARY 2016

DOI: 10.13140/RG.2.1.4091.4009

READS

21

3 AUTHORS, INCLUDING:



Salman Ahmed

University of Karachi

78 PUBLICATIONS 83 CITATIONS

[SEE PROFILE](#)



Zafar Alam Mahmood

COLORCON LIMITED - ENGLAND


64 PUBLICATIONS 71 CITATIONS

[SEE PROFILE](#)

4TH INTERNATIONAL CONFERENCE ON ENVIRONMENTAL HORIZON

Valuing and Conserving Nature

8-10 January 2016



PBS-01

TOTAL, INSOLUBLE AND SOLUBLE DIETARY FIBER CONTENTS OF THREE PAKISTANI LEGUME FLOURS

¹*Salman Ahmed, Muhammad Mohtasheemul Hasan¹ and

²Zafar Alam Mahmood


**¹ Department of Pharmacognosy, Faculty of Pharmacy, University of Karachi,
Karachi-75270, Pakistan.**

**² Colorcon Limited – UK, Flagship House, Victory Way, United Kingdom
E-mail: *salmanahmed@uok.edu.pk**

ABSTRACT

Current study was conducted to determine the insoluble, soluble and total dietary fiber content of *Macrotyloma uniflorum* (Lam.) Verdc., *Phaseolus lunatus* Linn., and *Phaseolus vulgaris* Linn., legume flours by using AOAC enzymatic - gravimetric analysis. The results showed that *M. uniflorum*, *P. lunatus* and *P. vulgaris* legume flours contained 155.91, 120.01 and 90.80 mg/g insoluble dietary fiber, respectively. Whereas, soluble dietary fiber was not found in any tested sample.

166



JOINTLY ORGANIZED BY :

Department of Chemistry & International Center for Chemical & Biological Sciences, University of Karachi